

Search for Leptoquarks with ATLAS at the LHC

Research Team

Th. D. Papadopoulou (P.I.), *Professor, National Technical University of Athens, Greece*

N. Ch. Benekos (*Researcher*), *Max-Planck-Institut fur Physik, Munich, Germany*

V. A. Mitsou (*Researcher*), *Instituto de Física Corpuscular, Valencia, Spain*

I. Panagoulas (*Research Student*), *National Technical University of Athens, Greece*

Introduction

Among possible new particles in physics beyond the Standard Model, leptoquarks (LQs) are an interesting category of exotic colour triplets with couplings to quarks and leptons. They are a generic prediction of Grand Unified Theories [1], of composite models [2], of technicolor schemes [3], of superstring-inspired E_6 models [4], and of supersymmetry with R -parity violation [5].

Leptoquarks are colour triplets which couple to quarks and leptons via a Yukawa-type coupling, λ , conserving the baryon and lepton numbers. In the model of Buchmüller, Rückl and Wyler (BRW) [6], leptoquarks couple to a single generation of Standard Model (SM) fermions via chiral Yukawa couplings which are invariant under the $SU(3)_C \times SU(2)_L \times U(1)_Y$ symmetry group. Inter-generational mixing is not allowed since neither flavour-changing neutral currents nor lepton flavour violation, induced by this mixing, have been observed so far. Moreover, leptoquarks coupling to both left- and right-handed electrons would mediate rare decays [7], hence leptoquarks couplings are assumed to be chiral.

There exist 14 species of leptoquarks, differing by their spin (scalars or vectors), fermion number $F=3B+L$, isospin and chirality of the coupling. They have fractional electric charge ($\pm 5/3$, $\pm 4/3$, $\pm 2/3$ and $\pm 1/3$) and decay into a charged or neutral lepton and a quark, according to the decay modes $LQ \rightarrow \ell q$ and $LQ \rightarrow \nu q$. In the most commonly used notation, $F=0$ scalar (S) and vector (V) species are denoted as S^L , $S_{1/2}^R$, $\tilde{S}_{1/2}$, V_0^L , V_0^R , \tilde{V}_0 and V_1 , while the $F=2$ species are S_0^L , S_0^R , \tilde{S}_0 , S_1 , $V_{1/2}^L$, $V_{1/2}^R$ and $\tilde{V}_{1/2}$. Decay widths for scalar and vector LQs of mass M_{LQ} are given by $\lambda^2 M_{LQ}/16\pi$ and $\lambda^2 M_{LQ}/24\pi$, respectively. The leptoquark branching fractions are predicted by the BRW model and take the values 1, 0.5 and 0.

In the following we consider pair production of scalar leptoquarks, which proceeds through gg fusion and $q\bar{q}$ annihilation via the diagrams shown in Figure 1. The production cross section is only slightly dependent on λ , since it is dominated by gg fusion which does not involve the ℓ -q-LQ coupling.




Figure 1. Leading-order diagrams of leptoquark pair production at LHC.

Furthermore, the cross section is not sensitive to the quark and lepton flavours to which the LQ couples, implying that all species are produced with the same rate. More information on LQ production at hadron colliders can be found in Ref. [8].

Event and detector simulation

In this study both the signal and the background events are generated with the event generator PYTHIA 6.2 [9], which provides a leading-order calculation of the LQ production cross section. The CTEQ5M set is used for the parton distribution functions and initial- and final-state radiation is switched on. As far as the Yukawa coupling is concerned, we set $\lambda = \sqrt{4\pi\alpha_{em}} \approx 0.3$, leading to a total decay width of $\Gamma \sim 2$ GeV for $M_{LQ} = 1$ TeV. This width is overwhelmed by the calorimeter resolution precluding any measurement of it, yet it allows one to treat the leptoquark as a resonance, i.e. it decays before fragmentation and only the products of its decay are observed in the detector. Hence the results obtained in this study are insensitive to λ as long as its value is consistent with resonant LQs, i.e. $\lambda \geq 10^{-6}$ for $M_{LQ} \geq 1$ TeV.

The performance of the ATLAS detector [10] was simulated with the fast simulation package ATLFast [11]. This simulation includes, in a parameterized way, the main aspects related to the detector response: jet reconstruction in the calorimeters, momentum/energy smearing for leptons and photons, reconstruction of missing transverse energy and charged particles. It is tuned to reproduce as well as possible the expected ATLAS performance [12].

Leptons plus jets channel ($\ell\ell jj$)

We have considered pair production of scalar leptoquarks of the first two generations. Both LQs are assumed to decay to a charged lepton and a quark, providing a topology with two high- p_T leptons and two high- E_T jets. Potential bounds in LQ mass are obtained for leptoquarks with branching ratio $BR(LQ \rightarrow \ell q) \equiv \beta$ equal to 1 or 0.5.

The background processes considered are characterized by the presence of two isolated high- p_T leptons and at least two jets in the final state. Before applying any cuts the main background source is the QCD processes. Nevertheless this is eliminated completely after the requirement for two isolated high- p_T leptons and in particular after selecting events with high lepton-jet reconstructed invariant mass, $m_{\ell j}$. The Drell-Yan lepton pair production, $q\bar{q} \rightarrow \gamma^*/Z \rightarrow \ell^+\ell^-$, is another possible background source but not as

important for the LHC (pp collider) as it is for Tevatron ($p\bar{p}$ collider). It can be eliminated in a similar manner after the high- m_{ℓ_j} cut is applied.

The cross sections of the surviving background processes are given in Table 1. The production of two leptons and two jets in the final state has been forced in PYTHIA. For processes with large cross section, such as Z+jet and $t\bar{t}$, the generation production has been divided in several \hat{p}_T intervals.

Processes	$\sigma \times BR$ (pb)	# events	# generated events
Z+jet ($\ell\ell jj$), $\hat{p}_T > 20$ GeV	1380	$4.14 \cdot 10^7$	$2.8 \cdot 10^6$
$t\bar{t}$ ($lvilvi$)	11	$3.3 \cdot 10^5$	$5 \cdot 10^5$
ZZ ($\ell\ell ii$)	1.2	$3.6 \cdot 10^4$	$2 \cdot 10^5$
ZW ($\ell\ell ij$)	1.2	$3.6 \cdot 10^4$	$2 \cdot 10^5$
WW ($lvlv$)	3.3	10^5	10^5

Table 1. Cross section, expected number of events for $L=30 \text{ fb}^{-1}$ and number of generated events of background processes for the LQ $LQ \rightarrow \ell\ell jj$ channel.

A first level of cuts is applied, requiring exactly two same-flavour, opposite-sign leptons with $p_T > 30$ GeV and $|\eta| < 2.5$, and at least two jets with $E_T > 30$ GeV and $|\eta| < 5$, for both signal and background, in order to identify the kinematic variables sensitive to the signal. The selection criteria have been optimized so that a significance over 5σ is achieved for the maximum possible leptoquark mass, retaining at the same time at least ten signal events for an integrated luminosity of 30 fb^{-1} .

The final selection cuts for both $eejj$ and $\mu\mu jj$ channels are the following: two same-flavour, opposite-sign leptons with $p_T > 100$ GeV and $|\eta| < 2.5$; at least two jets with $E_T > 70$ GeV and $|\eta| < 5$; invariant mass of the two leptons $m_{\ell\ell} > 180$ GeV, to exclude events from the $Z \rightarrow \ell\ell$ peak; $E_T^{\text{miss}} < 70$ GeV, to suppress the $t\bar{t}$ background; sum of transverse energy deposited in the calorimeters $\sum E_T^{\text{calo}} > 570$ GeV; ratio $E_T^{\text{miss}} / \sum E_T^{\text{calo}} < 0.05$ and a mass window $|m_{\ell_j} - M_{LQ}| < 100$ GeV. In the m_{ℓ_j} distribution, only the two leading jets and the two leptons are taken into account in the mass reconstruction. Moreover, between the two possible combinations, the one providing the minimum difference between the obtained values of m_{ℓ_j} is chosen.

The obtained m_{ℓ_j} distributions for the signal and the background, after applying the final cuts, are shown in Figure 2 for various leptoquark masses and for the $eejj$ channel. The cross section, the corresponding numbers of events in the mass window, as well as the significance achieved, are listed in Table 2 for a branching ratio of $\beta = 1$. The distributions are normalized to an integrated luminosity of 30 fb^{-1} , that is expected to be collected after three years of LHC running at low luminosity.

Signal can be clearly observed over the background for up to $M_{LQ} \approx 1.3$ TeV. For larger LQ masses, although the analysis is practically background-free, the low statistics preclude any potential observation of signal. Production cross section for the second generation ($\mu\mu jj$ channel), is about 2% lower, however, similar results are obtained.




Figure 2. m_{ej} distributions for background (light green line) and signal plus background (dark blue line), for various values of LQ mass and for an integrated luminosity of 30 fb^{-1} .

M_{LQ} (TeV)	σ (fb)	Signal	Background	S/\sqrt{B}
1.0	4.96	98.5	2.84	58
1.2	1.33	22.0	2.43	14
1.3	0.713	12.8	1.44	11
1.5	0.223	3.62	0.376	5.9

Table 2. Signal (1st generation) cross section, number of signal and background events and significance for the $eejj$ channel, for various values of LQ mass and for $L=30 \text{ fb}^{-1}$.

The leptoquarks studied so far decay with $\beta=1$ and correspond to the following particles: $S_0^R(-1/3)$, $\tilde{S}_0(-4/3)$, $S_{1/2}^L(-5/3)$, $S_{1/2}^R(-5/3)$, $S_{1/2}^R(-2/3)$, $\tilde{S}_{1/2}(-2/3)$, $S_1(-4/3)$. In the case of leptoquarks with $\beta=0.5$, i.e. $S_0^L(-1/3)$, $S_1(-1/3)$, the previous results become less stringent; leptoquarks are observable for $M_{LQ} \leq 1 \text{ TeV}$ with a significance of $S/\sqrt{B} \approx 15$ at $M_{LQ}=1 \text{ TeV}$.

Missing energy plus jets channel ($\nu\nu jj$)

In the $LQ LQ \rightarrow \nu\nu qq$ channel, both LQs are assumed to decay to a neutrino and a quark, providing a topology with two high- E_T jets and large missing transverse energy, E_T^{miss} . For the first two generations, the (huge) Z +jet ($\nu\nu jj$) background is irreducible making thus the analysis unfeasible. In contrast, for a third-generation leptoquark coupled to $\nu_\tau b$, the b-tagging capabilities of the ATLAS detector allow the signal to be disentangled from this background. Species with this coupling are studied in the following. The branching ratio of the decay $BR(LQ \rightarrow \nu_\tau b) = 1 - \beta$ is taken to be equal to 1 or 0.5.

Besides the Z +jet ($\nu\nu jj$) production, main background sources are the $t\bar{t}$ ($lvb\bar{b}lvb$), ZZ ($\nu\nu bb$) and ZW ($bb\bar{b}v$) processes. All other processes, listed in Table 3, are eliminated by the b-tagging and the lepton veto.

Processes	$\sigma \times BR$ (pb)	# events	# generated events
Z+jet (νjj)	22000	$6.6 \cdot 10^8$	$7.5 \cdot 10^5$
W+jet ($\ell \nu bb$)	28400	$8.5 \cdot 10^8$	$1.5 \cdot 10^5$
$t\bar{t}$ ($\ell \nu b \ell \nu b$)	51.6	$1.5 \cdot 10^6$	$5 \cdot 10^5$
ZZ ($\nu \nu bb$)	0.6	$1.8 \cdot 10^4$	10^5
ZW ($bb \ell \nu$)	1.3	$4 \cdot 10^4$	10^5
ZW ($\nu \nu jj$)	3.6	10^5	10^5
WW ($\ell \nu bb$)	30.5	$9 \cdot 10^5$	10^5

Table 3. Cross section, expected number of events for $L=30 \text{ fb}^{-1}$ and number of generated events of background processes for the LQ $LQ \rightarrow \nu bb$ channel.

Since in this channel leptoquarks decay to a τ -neutrino and a b-quark, producing two undetectable particles, the leptoquark mass is not reconstructed, and therefore only an excess of events can be observed. The different topology between signal and background events is taken into account by means of jet separation angles. Nevertheless, the signal selection can be improved if event shape variables, such as sphericity and aplanarity, are employed.

The final cuts applied to the event samples are the following: at least two jets tagged as originating from b-quarks with $E_T > 70 \text{ GeV}$ and $|\eta| < 5$; missing transverse energy $E_T^{\text{miss}} > 400 \text{ GeV}$; veto on isolated leptons for suppression of W+jets and $t\bar{t}$ background; and $m_{jj} > 180 \text{ GeV}$. In addition, since the two leptoquarks are produced back-to-back, the angular distribution of the final states should be constrained. Hence the azimuthal angle between the two leading jets is required to be $30^\circ < \Delta\phi_{j-j} < 150^\circ$, and the azimuthal angle between each of the two leading jets and the missing transverse momentum is selected to be $\Delta\phi(j-p_T^{\text{miss}}) > 60^\circ$.

As inferred from Table 4, for $\beta=0$ ($\tilde{S}_{1/2}(1/3)$) leptoquark observation is feasible for LQ masses of up to $\sim 1.3 \text{ TeV}$. The corresponding exclusion limit at 95% confidence level that this analysis can set is $M_{LQ} \leq 1.5 \text{ TeV}$. Moreover, if $\beta=0.5$ ($S_0^L(-1/3)$, $S_1^L(-1/3)$) — in which case LQs also couple to a τ -lepton and a top quark — the ATLAS reach using this channel reduces to $M_{LQ} \approx 1 \text{ TeV}$ with a significance of ~ 10 for $M_{LQ}=1 \text{ TeV}$.

M_{LQ} (TeV)	σ (fb)	Signal	Background	S/\sqrt{B}
1.0	4.84	70.7	3.4	38
1.2	1.28	21.3	3.4	12
1.3	0.68	12.1	3.4	6.5
1.5	0.21	3.9	3.4	2.1

Table 4. Signal (3rd generation) cross section, number of signal and background events and significance for the νbb channel, for various values of LQ mass and for $L=30 \text{ fb}^{-1}$.

Conclusions

The existence of TeV-scale scalar leptoquarks can be probed with the ATLAS experiment at the LHC for all fermion generations in pair production channels, extending the accessible LQ mass range by an order of magnitude compared to currently

running collider experiments. In the LQ $LQ \rightarrow \ell^+ \ell^- q \bar{q}$ mode the observation of a first- or second-generation leptoquark is feasible for up to $M_{LQ} \approx 1.3$ TeV ($M_{LQ} \approx 1$ TeV) assuming $\beta=1$ ($\beta=0.5$) with an integrated luminosity of 30 fb^{-1} . In the LQ $LQ \rightarrow \nu_\tau \nu_\tau b \bar{b}$ channel, on the other hand, third-generation LQs coupling to a τ -neutrino and a b-quark (b-anti-quark) with $\beta=0.5$ ($\beta=0$) will be equally probed for masses of up to 1 TeV (1.3 TeV).

It is emphasized that this analysis treats leptoquarks from a purely phenomenological point of view, i.e. the method is independent of the underlying theory. To this respect, it concerns any particle decaying to a lepton and a quark, e.g. squarks in some R -parity violating supersymmetric scenarios where lepton number is not conserved.

The results presented here are not meant to be considered as final, but rather represent an approximate yet robust idea of the ATLAS potential as far as leptoquark discovery is concerned. Further studies with GEANT-based simulated events will shed light on the actual profile of the signal events as seen from the ATLAS detector, possibly allowing more elaborate selection criteria to be imposed. Moreover, if next-to-leading order calculation [13] was employed for the LQ production cross section, the reach is expected to be enhanced. Study of other decay modes such as the ‘mixed’ LQ $LQ \rightarrow \ell q \nu q'$ channel will also allow to improve the reach.

References

1. J. C. Pati and A. Salam, *Phys. Rev. D* **10**, 275, 1974; H. Georgi and S. L. Glashow, *Phys. Rev. Lett.* **32**, 438, 1974
2. B. Schrempp and F. Schrempp, *Phys. Lett. B* **153**, 101, 1985; W. Buchmüller, *Acta Phys. Austriaca Suppl.* **27**, 517, 1985; W. Buchmüller and D. Wyler, *Phys. Lett. B* **177**, 377, 1986
3. S. Dimopoulos and L. Susskind, *Nucl. Phys. B* **155**, 237, 1979; S. Dimopoulos, *Nucl. Phys. B* **168**, 69, 1980; E. Eichten and K. Lane, *Phys. Lett. B* **90**, 125, 1980
4. V. D. Angelopoulos et al., *Nucl. Phys. B* **292**, 59, 1987; J. L. Hewett and T. G. Rizzo, *Phys. Rept.* **183**, 193, 1989
5. A. F. Žarnecki, *Eur. Phys. J. C* **17**, 695, 2000 [arXiv:hep-ph/0003271]
6. W. Buchmüller, R. Rückl and D. Wyler, *Phys. Lett. B* **191**, 442, 1987 [Erratum-ibid. *B* **448**, 320, 1999]
7. S. Davidson, D. C. Bailey and B. A. Campbell, *Z. Phys. C* **61**, 613, 1994 [arXiv:hep-ph/9309310]
8. J. L. Hewett and S. Pakvasa, *Phys. Rev. D* **37**, 3165, 1988; J. Blümlein, E. Boos and A. Kryukov, *Z. Phys. C* **76**, 137, 1997 [arXiv:hep-ph/9610408]
9. T. Sjöstrand et al., *Comput. Phys. Commun.* **135**, 238, 2001 [arXiv:hep-ph/0010017]
10. ATLAS Collaboration, *ATLAS Technical Proposal*, CERN/LHCC/94-43, 1994
11. E. Richter-Was, D. Froidevaux and L. Poggioli, *ATLFAST 2.0: a fast simulation package for ATLAS*, ATLAS Internal Note, ATL-PHYS-98-131, 1998
12. ATLAS Collaboration, *Detector and Physics Performance Technical Design Report* vol. I, CERN/LHCC/99-14, 1999
13. M. Krämer, T. Plehn, M. Spira and P. M. Zerwas, *Phys. Rev. Lett.* **79**, 341, 1997 [arXiv:hep-ph/9704322]; arXiv:hep-ph/04110